

A RADON-NIKODYM THEOREM FOR COMPLETELY MULTI-POSITIVE LINEAR MAPS AND ITS APPLICATIONS

MARIA JOIȚA

February 2, 2008

Abstract

A completely n -positive linear map from a locally C^* -algebra A to another locally C^* -algebra B is an $n \times n$ matrix whose elements are continuous linear maps from A to B and which verifies the condition of completely positivity. In this paper we prove a Radon-Nikodym type theorem for strict completely n -positive linear maps which describes the order relation on the set of all strict completely n -positive linear maps from a locally C^* -algebra A to a C^* -algebra B , in terms of a self-dual Hilbert C^* -module structure induced by each strict completely n -positive linear map. As applications of this result we characterize the pure completely n -positive linear maps from A to B and the extreme elements in the set of all identity preserving completely n -positive linear maps from A to B . Also we determine a certain class of extreme elements in the set of all identity preserving completely positive linear maps from A to $M_n(B)$.

MSC: 46L05; 46L08

1 Introduction and preliminaries

The concept of matricial order plays an important role to understand the infinite -dimensional non-commutative structure of operator algebras. Completely positive maps, as the natural ordering attached to this structure have been studied extensively [1, 2, 5, 4, 7, 8, 9, 10, 11, 15, 16, 18, 19].

Given a C^* -algebra A we denote by $M_n(A)$ the C^* -algebra of all $n \times n$ matrices with elements in A .

DEFINITION 1.1 *A completely positive map from a C^* -algebra A to another C^* -algebra B is a linear map $\rho : A \rightarrow B$ such that the linear map $\rho_n :$*

$M_n(A) \rightarrow M_n(B)$ defined by

$$\rho_n \left([a_{ij}]_{i,j=1}^n \right) = [\rho(a_{ij})]_{i,j=1}^n$$

is positive for each positive integer n .

In 1955, Stinespring [18] showed that a completely positive linear map ρ from a C^* -algebra A to $L(H)$, the C^* -algebra of all bounded linear operators on a Hilbert space H , induces a representation Φ_ρ of A on another Hilbert space H_ρ . Moreover,

$$\rho(a) = V_\rho^* \Phi_\rho(a) V_\rho$$

for all $a \in A$ and for some bounded linear operator V_ρ from H to H_ρ .

In 1969, Arveson [1] proved a Radon-Nikodym type theorem which gives a description of the order relation in the set of all completely positive linear maps from A to $L(H)$ in terms of the Stinespring representation associated with each completely positive linear map. The gist of the proof of this result is the fact that any bounded linear operator on a Hilbert space is adjointable.

Hilbert C^* -modules are generalizations of Hilbert spaces by allowing the inner-product to take values in a C^* -algebra rather than in the field of complex numbers.

DEFINITION 1.2 *A pre-Hilbert A -module is a complex vector space E which is also a right A -module, compatible with the complex algebra structure, equipped with an A -valued inner product $\langle \cdot, \cdot \rangle : E \times E \rightarrow A$ which is $\mathbb{C}$ - and A -linear in its second variable and satisfies the following relations:*

1. $\langle \xi, \eta \rangle^* = \langle \eta, \xi \rangle$ for every $\xi, \eta \in E$;
2. $\langle \xi, \xi \rangle \geq 0$ for every $\xi \in E$;
3. $\langle \xi, \xi \rangle = 0$ if and only if $\xi = 0$.

We say that E is a Hilbert A -module if E is complete with respect to the topology determined by the norm $\|\cdot\|$ given by $\|\xi\| = \sqrt{\|\langle \xi, \xi \rangle\|}$.

A C^* -algebra A is a Hilbert C^* -module over A with the inner-product defined by $\langle a, b \rangle = a^*b$ for a and b in A .

Given two Hilbert A -modules E and F , the Banach space of all bounded module homomorphisms from E to F is denoted by $\mathcal{B}_A(E, F)$. The subset of $\mathcal{B}_A(E, F)$ consisting of all adjointable module homomorphisms from E to F (that is, $T \in \mathcal{B}_A(E, F)$ such that there is $T^* \in \mathcal{B}_A(F, E)$ satisfying

$\langle \eta, T\xi \rangle = \langle T^*\eta, \xi \rangle$ for all $\xi \in E$ and for all $\eta \in F$) is denoted by $\mathcal{L}_A(E, F)$. We will write $\mathcal{B}_A(E)$ for $\mathcal{B}_A(E, E)$ and $\mathcal{L}_A(E)$ for $\mathcal{L}_A(E, E)$.

In general, $\mathcal{L}_A(E, F) \neq \mathcal{B}_A(E, F)$. So the theory of Hilbert C^* -modules is different from the theory of Hilbert spaces.

The Banach space $E^\#$ of all bounded module homomorphisms from E to A becomes a right A -module with the action of A on $E^\#$ defined by $(aT)(\xi) = a^*(T\xi)$ for $a \in A$, $T \in E^\#$ and $\xi \in E$. We say that E is self-dual if $E^\# = E$ as right A -modules.

If E and F are self-dual, then $\mathcal{B}_A(E, F) = \mathcal{L}_A(E, F)$ [16, Proposition 3.4].

Suppose that A is a W^* -algebra. Then the A -valued inner-product on E extends to an A -valued inner-product on $E^\#$ and in this way $E^\#$ becomes a self-dual Hilbert A -module [16, Theorem 3.2]. Moreover, any bounded module homomorphism T from E to F extends uniquely to a bounded homomorphism $\tilde{T}$ from $E^\#$ to $F^\#$ [16, Proposition 3.6].

A representation of a C^* -algebra A on a Hilbert C^* -module E over a C^* -algebra B is a $*$ -morphism Φ from A to $\mathcal{L}_B(E)$.

Paschke [16] showed that a completely positive map from a unital C^* -algebra A to another unital C^* -algebra B induces a representation of A on a Hilbert B -module which generalizes the GNS construction and he extended the Arveson's results for completely positive maps from a unital C^* -algebra A to a W^* -algebra B .

In 1996, Tsui [19] proved a Radon-Nikodym type theorem for completely positive maps between unital C^* -algebras and using this theorem he obtained characterizations of the pure elements and the extreme points in the set of all identity preserving completely positive maps from a unital C^* -algebra A to another unital C^* -algebra B in terms of a self-dual Hilbert module structure induced by each completely positive map. To prove these facts he used the following construction.

CONSTRUCTION 1.3 ([15, 16, 19]) *Let E be a Hilbert C^* -module over a C^* -algebra B . The algebraic tensor product $E \otimes_{alg} B^{**}$, where B^{**} is the enveloping W^* -algebra of B , becomes a right B^{**} -module if we define $(\xi \otimes b)c = \xi \otimes bc$, for $\xi \in E$, and $b, c \in B^{**}$.*

*The map $[\cdot, \cdot] : (E \otimes_{alg} B^{**}) \times (E \otimes_{alg} B^{**}) \rightarrow B^{**}$ defined by*

$$\left[\sum_{i=1}^n \xi_i \otimes b_i, \sum_{j=1}^m \eta_j \otimes c_j \right] = \sum_{i=1}^n \sum_{j=1}^m b_i^* \langle \xi_i, \eta_j \rangle c_j$$

is a B^{**} -valued inner-product on $E \otimes_{\text{alg}} B^{**}$ and the quotient module $E \otimes_{\text{alg}} B^{**}/N_E$, where $N_E = \{\zeta \in E \otimes_{\text{alg}} B^{**}; [\zeta, \zeta] = 0\}$, becomes a pre-Hilbert B^{**} -module. The Hilbert C^* -module $\overline{E \otimes_{\text{alg}} B^{**}/N_E}$ obtained by the completion of $E \otimes_{\text{alg}} B^{**}/N_E$ with respect to the norm induced by the inner product $[\cdot, \cdot]$ is called the extension of E by the C^* -algebra B^{**} . Moreover, E can be regarded as a B -submodule of $\overline{E \otimes_{\text{alg}} B^{**}/N_E}$, since the map $\xi \mapsto \xi \otimes 1 + N_E$ from E to $\overline{E \otimes_{\text{alg}} B^{**}/N_E}$ is an isometric inclusion.

The self-dual Hilbert B^{**} -module $\left(\overline{E \otimes_{\text{alg}} B^{**}/N_E}\right)^\#$ is denoted by $\tilde{E}$, and we can consider E as embedded in $\tilde{E}$ without making distinction.

Let $T \in \mathcal{B}_B(E, F)$. For $b_1, \dots, b_m \in B^{**}$ and $\xi_1, \dots, \xi_m$ in E we denote by b the element in $(B^{**})^m$ whose components are $b_1, \dots, b_m$, by X the matrix in $M_n(B^{**})$ whose the (i, j) -entry is $\langle \xi_i, \xi_j \rangle$ and by X_T the matrix in $M_n(B^{**})$ whose the (i, j) -entry is $\langle T\xi_i, T\xi_j \rangle$. By Lemma 4.2 in [14], $0 \leq X_T \leq \|T\| X$. Identifying $M_n(B^{**})$ with $L_{B^{**}}((B^{**})^n)$, we have

$$\begin{aligned} \left[\sum_{i=1}^m T\xi_i \otimes b_i, \sum_{i=1}^m T\xi_i \otimes b_i \right] &= \sum_{i,j=1}^m b_i^* \langle T\xi_i, T\xi_j \rangle b_j = \langle b, X_T b \rangle \\ &\leq \|T\| \langle b, X b \rangle = \|T\| \left[\sum_{i=1}^m \xi_i \otimes b_i, \sum_{i=1}^m \xi_i \otimes b_i \right]. \end{aligned}$$

Therefore T extends uniquely to a bounded module homomorphism $\hat{T}$ from $\overline{E \otimes_{\text{alg}} B^{**}/N_E}$ to $\overline{F \otimes_{\text{alg}} B^{**}/N_F}$ such that

$$\hat{T} \left(\sum_{i=1}^m \xi_i \otimes b_i \right) = \sum_{i=1}^m T\xi_i \otimes b_i$$

and by Proposition 3.6 in [16], this extends uniquely to a bounded module homomorphism $\tilde{T}$ from $\tilde{E}$ to $\tilde{F}$ such that $\|T\| = \|\tilde{T}\|$.

REMARK 1.4 Any element $T \in \mathcal{B}_B(E, F)$ extends uniquely to an element $\tilde{T} \in \mathcal{B}_{B^{**}}(\tilde{E}, \tilde{F})$ such that $\|T\| = \|\tilde{T}\|$. Moreover, $\widetilde{TS} = \tilde{T}\tilde{S}$ for all $T \in \mathcal{B}_B(E, F)$ and $S \in \mathcal{B}_B(F, E)$, and if $T \in \mathcal{L}(E, F)$, then $\widetilde{T^*} = \tilde{T}^*$.

REMARK 1.5 Let $T \in \mathcal{B}_B(E, E^\#)$. We extend T to an element $\overline{T} \in \mathcal{B}_B((E \otimes_{alg} B^{**})/N_E, \tilde{E})$ by

$$\left[\overline{T} \left(\sum_{i=1}^n \xi_i \otimes b_i \right), \sum_{j=1}^m \eta_j \otimes c_j \right] = \sum_{i=1}^n \sum_{j=1}^m b_i^* [T\xi_i, \eta_j] c_j$$

and then extend it again to an element $\tilde{T} \in \mathcal{B}_B(\tilde{E})$ such that $\|T\| = \|\tilde{T}\|$ [16, Proposition 3.6].

REMARK 1.6 A representation Φ of a C^* -algebra A on a Hilbert C^* -module E over a C^* -algebra B induces a representation $\tilde{\Phi}$ of A on $\tilde{E}$ defined by $\tilde{\Phi}(a) = \widetilde{\Phi(a)}$ for all $a \in A$.

REMARK 1.7 Any completely positive linear map ρ from A to B induces a representation $\tilde{\Phi}_\rho$ of A on a self-dual Hilbert B^{**} -module $\tilde{E}_\rho$.

Locally C^* -algebras are generalizations of C^* -algebras. Instead of being given by a single norm, the topology on a locally C^* -algebra is defined by a directed family of C^* -seminorms. In fact a locally C^* -algebra is an inverse limit of C^* -algebras.

DEFINITION 1.8 A locally C^* -algebra is a complete complex Hausdorff topological $*$ -algebra A whose topology is determined by its continuous C^* -seminorms in the sense that the net $\{a_i\}_{i \in I}$ converges to 0 if and only if the net $\{p(a_i)\}_{i \in I}$ converges to 0 for all continuous C^* -seminorm p on A .

If A is a locally C^* -algebra and $S(A)$ is the set of all continuous C^* -seminorms on A , then for each $p \in S(A)$, $A_p = A/\ker p$ is a C^* -algebra in the norm induced by p . The canonical map from A onto A_p is denoted by π_p for each $p \in S(A)$. For $p, q \in S(A)$ with $q \leq p$ there is a unique morphism of C^* -algebras π_{pq} from A_p onto A_q such that $\pi_{pq}(\pi_p(a)) = \pi_q(a)$ for all $a \in A$. Moreover, $\{A_p; \pi_{pq}\}_{p, q \in S(A), p \geq q}$ is an inverse system of C^* -algebras, and A can be identified with $\varprojlim_p A_p$. Clearly, any C^* -algebra is a locally C^* -algebra.

The terminology "locally C^* -algebra" is due to Inoue [6]. In the literature, locally C^* -algebras have been given different name such as b^* -algebras, m -convex- C^* -algebras, LMC^* -algebras [3] or pro- C^* -algebras [17]. Such important concepts as Hilbert C^* -modules, adjointable operators, (completely) positive linear maps, (completely) multi-positive linear maps can be

defined with obvious modifications in the framework of locally C^* -algebras and many results from the theory of C^* -algebras are still valid. The proofs are not always straightforward. Thus, in [3] it is proved that a continuous positive functional ρ on a locally C^* -algebra A induces a representation of A on a Hilbert space H which extends the GNS construction, and moreover, the representation of A induced by ρ is irreducible if and only if ρ is pure. In [2], Bhatt and Karia extend the Stinespring construction for completely positive linear maps from a locally C^* -algebra A to $L(H)$.

If A is a locally C^* -algebra, then the set $M_n(A)$ of all $n \times n$ matrices over A with the algebraic operations and the topology obtained by replying it as a direct sum of n^2 copies of A is a locally C^* -algebra.

DEFINITION 1.9 ([4], [9]). *A completely n -positive map from a locally C^* -algebra A to another locally C^* -algebra B is an $n \times n$ matrix $[\rho_{ij}]_{i,j=1}^n$ whose elements are continuous linear maps from A to B such that the map ρ from $M_n(A)$ to $M_n(B)$ defined by*

$$\rho \left([a_{ij}]_{i,j=1}^n \right) = [\rho_{ij}(a_{ij})]_{i,j=1}^n$$

is completely positive.

DEFINITION 1.10 ([9]). *A completely n -positive map $[\rho_{ij}]_{i,j=1}^n$ from A to $\mathcal{L}_B(E)$, where E is a Hilbert module over a C^* -algebra B is strict if for some approximate unit $\{e_\lambda\}_{\lambda \in \Lambda}$ for A , the nets $\{\rho_{ii}(e_\lambda)\}_{\lambda \in \Lambda}$, $i \in \{1, \dots, n\}$ are strictly Cauchy in $\mathcal{L}_B(E)$ (that is, the nets $\{\rho_{ii}(e_\lambda)\xi\}_{\lambda \in \Lambda}$, $i \in \{1, \dots, n\}$ are Cauchy in E for each $\xi \in E$).*

REMARK 1.11 *If A is unital or E is a Hilbert space, then any completely n -positive map from A to $\mathcal{L}(E)$ is strict.*

In [9], we extend the KSGNS (Kasparov, Stinespring, Gel'fand, Naimark, Segal) construction for strict completely multi-positive linear maps between locally C^* -algebras.

Theorem 1 ([9]). *Let A be a locally C^* -algebras, let E be a Hilbert module over a C^* -algebra B and let $\rho = [\rho_{ij}]_{i,j=1}^n$ be a strict completely n -positive map from A to $\mathcal{L}_B(E)$.*

1. *There is a representation Φ_ρ of A on a Hilbert B -module E_ρ and there are n elements $V_{\rho,1}, \dots, V_{\rho,n}$ in $\mathcal{L}_B(E, E_\rho)$ such that*

- (a) $\rho_{ij}(a) = V_{\rho,i}^* \Phi_\rho(a) V_{\rho,j}$ for all $a \in A$ and for all $i, j \in \{1, \dots, n\}$;
- (b) $\{\Phi_\rho(a) V_{\rho,i} x; a \in A, x \in E, 1 \leq i \leq n\}$ spans a dense subspace of E_ρ .

2. If Φ is another representation of A on a Hilbert B -module F and $W_1, \dots, W_n$ are n elements in $\mathcal{L}_B(E, F)$ such that

- (a) $\rho_{ij}(a) = W_i^* \Phi(a) W_j$ for all $a \in A$ and for all $i, j \in \{1, \dots, n\}$;
- (b) $\{\Phi(a) W_i x; a \in A, x \in E, 1 \leq i \leq n\}$ spans a dense subspace of F ,
there is a unitary operator $U \in \mathcal{L}_B(E_\rho, F)$ such that
 - i. $\Phi(a) U = U \Phi_\rho(a)$ for all $a \in A$; and
 - ii. $W_i = U V_{\rho,i}$ for all $i \in \{1, \dots, n\}$.

The $n+2$ tuple $(\Phi_\rho, E_\rho, V_{\rho,1}, \dots, V_{\rho,n})$ is called the KSGNS construction associated with ρ .

In [10], we prove a Radon-Nikodym type theorem for completely multi-positive linear maps from a locally C^* -algebra A to $L(H)$ and we characterize the pure elements and the extreme points in the set of all identity preserving completely multi-positive linear maps from A to $L(H)$ in terms of the representation of A induced by each completely multi-positive linear map. Also, we determine a certain class of extreme points in the set of all identity preserving completely positive linear maps from A to $M_n(L(H))$. In this talk, we will extend the results from [10] for completely multi-positive linear maps from a locally C^* -algebra A to a C^* -algebra B .

2 The Radon-Nikodym theorem for completely n -positive linear maps

Throughout this section, we assume that A is a locally C^* -algebra, B is a C^* -algebra and E is a Hilbert C^* -module over B . We will denote by $SCP_\infty^n(A, \mathcal{L}_B(E))$ the set of all strict completely n -positive linear maps from A to $\mathcal{L}_B(E)$ and by $CP_\infty(A, \mathcal{L}_B(E))$ the set of all completely positive linear maps from A to $\mathcal{L}_B(E)$.

PROPOSITION 2.1 ([4],[10]) *There is a bijection $\mathcal{S}$ from the set $CP_\infty^n(A, B)$ of all completely n -positive maps from A to B onto the set $CP_\infty(A, M_n(B))$ of all completely positive maps from A to $M_n(B)$ defined by*

$$\mathcal{S} \left([\rho_{ij}]_{i,j=1}^n \right) (a) = [\rho_{ij}(a)]_{i,j=1}^n \text{ for all } a \in A$$

which preserves the order relation.

For an element $T \in \mathcal{L}_{B^{**}}(\widetilde{E})$ we denote by $T|_E$ the restriction of the map T on E .

Let $\rho \in SCP_\infty^n(A, \mathcal{L}_B(E))$. We denote by $C(\rho)$ the C^* -subalgebra of $\mathcal{L}_{B^{**}}(\widetilde{E}_\rho)$ generated by $\{T \in \mathcal{L}_{B^{**}}(\widetilde{E}_\rho); T\widetilde{\Phi}_\rho(a) = \widetilde{\Phi}_\rho(a)T, \widetilde{V_{\rho,j}}^* T\widetilde{\Phi}_\rho(a)\widetilde{V_{\rho,i}}|_E \in \mathcal{L}_B(E) \text{ for all } a \in A \text{ and for all } i, j \in \{1, \dots, n\}\}$.

REMARK 2.2 If T is an element in $C(\rho)$, then $T|_{E_\rho} \in \mathcal{B}_B(E_\rho, E_\rho^\#)$, since

$$\langle T\widetilde{\Phi}_\rho(a)V_{\rho,j}\xi, \widetilde{\Phi}_\rho(b)V_{\rho,i}\eta \rangle = \langle T\widetilde{\Phi}_\rho(a)\widetilde{V_{\rho,j}}\xi, \widetilde{\Phi}_\rho(b)\widetilde{V_{\rho,i}}\eta \rangle \in B$$

for all $a, b \in A$, for all $\xi, \eta \in E$ and for all $i, j \in \{1, \dots, n\}$ and since $\{\widetilde{\Phi}_\rho(a)V_{\rho,i}\xi; a \in A, \xi \in E, 1 \leq i \leq n\}$ spans a dense submodule of E_ρ .

LEMMA 2.3 Let $T \in C(\rho)$. If T is positive, then the map ρ_T from $M_n(A)$ to $M_n(\mathcal{L}_B(E))$ defined by

$$\rho_T([a_{ij}]_{i,j=1}^n) = [\widetilde{V_{\rho,i}}^* T\widetilde{\Phi}_\rho(a_{ij})\widetilde{V_{\rho,j}}|_E]_{i,j=1}^n$$

is a strict completely n -positive linear map from A to $\mathcal{L}_B(E)$.

Proof. It is not difficult to see that ρ_T is an $n \times n$ matrix of continuous linear maps from A to $\mathcal{L}_B(E)$, the (i, j) -entry of the matrix ρ_T is the linear map $(\rho_T)_{ij}$ from A to $\mathcal{L}_B(E)$ defined by

$$(\rho_T)_{ij}(a) = \widetilde{V_{\rho,i}}^* T\widetilde{\Phi}_\rho(a)\widetilde{V_{\rho,j}}|_E.$$

Also it is not difficult to check that for all $a_1, \dots, a_m \in A$ and for all $T_1, \dots, T_m \in M_n(\mathcal{L}_B(E))$, we have

$$\begin{aligned} \sum_{k,l=1}^m T_l^* \mathcal{S}(\rho_T)(a_l^* a_k) T_k &= \left(\sum_{k,l=1}^m \widetilde{T_l}^* \widetilde{\mathcal{S}(\rho_T)}(a_l^* a_k) \widetilde{T_k} \right) \Big|_E \\ &= \left(\left(\sum_{l=1}^m M_{T_l^{\frac{1}{2}}}(a_l) V \widetilde{T_l} \right)^* \left(\sum_{l=1}^m M_{T_l^{\frac{1}{2}}}(a_l) V \widetilde{T_l} \right) \right) \Big|_E, \end{aligned}$$

$$\text{where } M_{T_l^{\frac{1}{2}}}(a) = \begin{bmatrix} T^{\frac{1}{2}}\widetilde{\Phi}_\rho(a) & \cdots & T^{\frac{1}{2}}\widetilde{\Phi}_\rho(a) \\ 0 & \cdots & 0 \\ \vdots & \cdots & \vdots \\ 0 & \cdots & 0 \end{bmatrix} \text{ and } V = \begin{bmatrix} \widetilde{V_{\rho,1}} & \cdots & 0 \\ \vdots & \cdots & \vdots \\ 0 & \cdots & \widetilde{V_{\rho,n}} \end{bmatrix}.$$

From this fact we conclude that $\mathcal{S}(\rho_T) \in CP_\infty(A, M_n(\mathcal{L}_B(E)))$ and by

Proposition 2.1, $\rho_T \in CP_\infty^n(A, \mathcal{L}_B(E))$. To show that $\rho_T \in SCP_\infty^n(A, \mathcal{L}_B(E))$, let $\{e_\lambda\}_{\lambda \in \Lambda}$ be an approximate unit for A , $\xi \in E$ and $i \in \{1, \dots, n\}$. Then

$$\begin{aligned} \|(\rho_T)_{ii}(e_\lambda)\xi - (\rho_T)_{ii}(e_\mu)\xi\| &= \left\| \widetilde{V_{\rho,i}}^* T \left(\widetilde{\Phi_\rho}(e_\lambda) - \widetilde{\Phi_\rho}(e_\mu) \right) \widetilde{V_{\rho,i}} \xi \right\| \\ &\leq \left\| \widetilde{V_{\rho,i}}^* T \right\| \left\| (\Phi_\rho(e_\lambda) - \Phi_\rho(e_\mu)) V_{\rho,i} \xi \right\|, \end{aligned}$$

and since $\{\Phi_\rho(e_\lambda) V_{\rho,i} \xi\}_{\lambda \in \Lambda}$ is a Cauchy net in E , the net $\{(\rho_T)_{ii}(e_\lambda)\}_{\lambda \in \Lambda}$ is strictly Cauchy. Therefore $\rho_T \in SCP_\infty^n(A, \mathcal{L}_B(E))$. ■

REMARK 2.4 1. If $I_{\widetilde{E_\rho}}$ is the identity map on $\widetilde{E_\rho}$, then $\rho_{I_{\widetilde{E_\rho}}} = \rho$.

2. If T_1 and T_2 are two positive elements in $C(\rho)$, then $\rho_{T_1+T_2} = \rho_{T_1} + \rho_{T_2}$.

3. If T is a positive element in $C(\rho)$ and α is a positive number, then $\rho_{\alpha T} = \alpha \rho_T$.

REMARK 2.5 Let T_1 and T_2 be two positive elements in $C(\rho)$. If $T_1 \leq T_2$, then, since

$$\begin{aligned} (\rho_{T_2} - \rho_{T_1}) \left([a_{ij}]_{i,j=1}^n \right) &= \left[\widetilde{V_{\rho,i}}^* (T_2 - T_1) \widetilde{\Phi_\rho}(a_{ij}) \widetilde{V_{\rho,j}} \right]_{i,j=1}^n \\ &= \rho_{T_2 - T_1} \left([a_{ij}]_{i,j=1}^n \right) \end{aligned}$$

for all $[a_{ij}]_{i,j=1}^n \in M_n(A)$, $\rho_{T_1} \leq \rho_{T_2}$.

Let $\rho \in SCP_\infty^n(A, \mathcal{L}_B(E))$. We denote by $[0, \rho]$ the set of all strict completely n -positive linear maps θ from A to $\mathcal{L}_B(E)$ such that $\theta \leq \rho$ (that is, $\rho - \theta \in SCP_\infty^n(A, \mathcal{L}_B(E))$) and by $[0, I]_\rho$ the set of all elements T in $C(\rho)$ such that $0 \leq T \leq I_{\widetilde{E_\rho}}$.

THEOREM 2.6 The map $T \rightarrow \rho_T$ from $[0, I]_\rho$ to $[0, \rho]$ is an affine order isomorphism.

Proof. By Lemma 2.3 and Remarks 2.4 and 2.5, the map $T \rightarrow \rho_T$ from $[0, I]_\rho$ to $[0, \rho]$ is well-defined and moreover, it is affine. To show that this map is injective, let $T \in [0, I]_\rho$ such that $\rho_T = 0$. Then $\widetilde{V_{\rho,i}}^* T \widetilde{\Phi_\rho}(a) \widetilde{V_{\rho,j}} \Big|_E = 0$ for all $a \in A$ and for all $i, j \in \{1, 2, \dots, n\}$, and so

$$\langle T \Phi_\rho(a) V_{\rho,j} \xi, \Phi_\rho(b) V_{\rho,j} \eta \rangle = 0$$

for all $a, b \in A$ for all $\xi, \eta \in E$ and for all $i, j \in \{1, \dots, n\}$. Taking into account that $\{\Phi_\rho(a)V_{\rho,i}\xi; a \in A, \xi \in E, 1 \leq i \leq n\}$ spans a dense submodule of E_ρ , from these facts, Remarks 2.2 and 1.5 and we conclude that $T = 0$.

Let $\theta \in [0, \rho]$. In the same way as in the proof of Lemma 3.4 in [10], we show that there is a bounded linear map W from E_ρ to E_θ such that

$$W(\Phi_\rho(a)V_{\rho,i}\xi) = \Phi_\theta(a)V_{\theta,i}\xi.$$

It is not difficult to check that W is a bounded module homomorphism such that $\|W\| \leq 1$, $WV_{\rho,i} = V_{\theta,i}$ for all $i \in \{1, \dots, n\}$ and $W\Phi_\rho(a) = \Phi_\theta(a)W$ for all $a \in A$. If $\widetilde{W}$ is the unique extension of W to a bounded module morphism from $\widetilde{E}_\rho$ to $\widetilde{E}_\theta$ with $\|\widetilde{W}\| = \left\| \widetilde{W} \right\|$, then clearly $0 \leq \widetilde{W}^* \widetilde{W} \leq I_{\widetilde{E}_\rho}$. Moreover, it is easy to check that $\widetilde{W}^* \widetilde{W} \Phi_\rho(a) = \Phi_\rho(a) \widetilde{W}^* \widetilde{W}$ for all $a \in A$, and since

$$\widetilde{V}_{\rho,i}^* \widetilde{W}^* \widetilde{W} \Phi_\rho(a) \widetilde{V}_{\rho,j} = \widetilde{V}_{\rho,i}^* \widetilde{W}^* \Phi_\theta(a) \widetilde{W} \widetilde{V}_{\rho,j} = \widetilde{V}_{\theta,i}^* \Phi_\theta(a) \widetilde{V}_{\theta,j}$$

for all $a \in A$ and for all $i, j \in \{1, \dots, n\}$, $\widetilde{W}^* \widetilde{W} \in [0, I]_\rho$. Let $T = \widetilde{W}^* \widetilde{W}$. Then clearly, $\theta = \rho_T$ and thus the map $T \rightarrow \rho_T$ from $[0, I]_\rho$ to $[0, \rho]$ is surjective. Therefore the map $T \rightarrow \rho_T$ is an affine isomorphism from $[0, I]_\rho$ onto $[0, \rho]$ which preserve the order relation. ■

3 Applications of the Radon-Nikodym theorem

Let A be a locally C^* -algebra, let B be a C^* -algebra and let E be a Hilbert C^* -module over B . A strict completely n -positive linear map ρ from A to $\mathcal{L}_B(E)$ is said to be pure if for every strict completely n -positive linear map θ from A to $\mathcal{L}_B(E)$ with $\theta \leq \rho$, there is a positive number α such that $\theta = \alpha\rho$.

PROPOSITION 3.1 *Let $\rho \in SCP_\infty^n(A, \mathcal{L}_B(E))$. Then ρ is pure if and only if $[0, I]_\rho = \{\alpha I_{\widetilde{E}_\rho}; 0 \leq \alpha \leq 1\}$.*

Proof. First we suppose that ρ is pure. Let $T \in [0, I]_\rho$. By Theorem 2.6, $\rho_T \in [0, \rho]$, and since ρ is pure, $\rho_T = \alpha\rho$ for some positive number. From this fact, Remark 2.4 and Theorem 2.6 we deduce that $T = \alpha I_{\widetilde{E}_\rho}$ for some $0 \leq \alpha \leq 1$.

Conversely, suppose that $[0, I]_\rho = \{\alpha I_{\widetilde{E}_\rho}; 0 \leq \alpha \leq 1\}$. Let $\theta \in SCP_\infty^n(A, \mathcal{L}_B(E))$ such that $\theta \leq \rho$. By Theorem 2.6, $\theta = \rho_T$ for some $T \in [0, I]_\rho$, and since $T = \alpha I_{\widetilde{E}_\rho}$ for some positive number α , $\theta = \alpha\rho$ and the proposition is proved. ■

COROLLARY 3.2 *A strict completely n -positive linear map ρ from A to $\mathcal{L}_B(E)$ is pure if and only if $C(\rho)$ consisting of the scalar multipliers of $I_{\widetilde{E_\rho}}$.*

We say that two strict completely n -positive linear maps ρ and θ from A to $\mathcal{L}_B(E)$ are unitarily equivalent if the representations of A induced by ρ respectively θ are unitarily equivalent.

The following proposition is a generalization of Proposition 4.3 in [10].

PROPOSITION 3.3 *Let A be a unital locally C^* -algebra, let B be a C^* -algebra, let E be a Hilbert B -module and let $\rho = [\rho_{ij}]_{i,j=1}^n \in CP_\infty^n(A, \mathcal{L}_B(E))$. If ρ_{ii} , $i \in \{1, \dots, n\}$ are unitarily equivalent pure unital completely positive linear maps from A to $\mathcal{L}_B(E)$ and for all $i, j \in \{1, \dots, n\}$ with $i \neq j$ there is a unitary element u_{ij} in A such that $\rho_{ij}(u_{ij}) = I_E$, then ρ is pure.*

Proof. Let $i, j \in \{1, \dots, n\}$ with $i \neq j$. From

$$\|\Phi_\rho(u_{ij})V_{\rho,j} - V_{\rho,i}\|^2 = \|V_{\rho,j}^*V_{\rho,j} - \rho_{ij}(u_{ij}) - (\rho_{ij}(u_{ij}))^* + V_{\rho,i}^*V_{\rho,i}\| = 0$$

we deduce that $\Phi_\rho(u_{ij})V_{\rho,j} = V_{\rho,i}$. Therefore the sets $\{\Phi_\rho(a)V_{\rho,i}\xi; a \in A, \xi \in E\}$ and $\{\Phi_\rho(a)V_{\rho,j}\xi; a \in A, \xi \in E\}$ generate the same Hilbert submodule of E_ρ , and since E_ρ is generated by $\{\Phi_\rho(a)V_{\rho,i}\xi; a \in A, \xi \in E, 1 \leq i \leq n\}$, this coincides with E_ρ .

Let $i \in \{1, \dots, n\}$ and let $(\Phi_{\rho_{ii}}, E_{\rho_{ii}}, V_{\rho_{ii}})$ be the KSGNS construction associated with ρ_{ii} . We will show that the representations Φ_ρ and $\Phi_{\rho_{ii}}$ of A are unitarily equivalent. Since $\{\Phi_\rho(a)V_{\rho,i}\xi; a \in A, \xi \in E\}$ spans a dense submodule of E_ρ , $\{\Phi_{\rho_{ii}}(a)V_{\rho_{ii}}\xi; a \in A, \xi \in E\}$ spans a dense submodule of $E_{\rho_{ii}}$ and

$$\begin{aligned} \langle \Phi_\rho(a)V_{\rho,i}\xi, \Phi_\rho(b)V_{\rho,i}\eta \rangle &= \langle \rho_{ii}(b^*a)\xi, \eta \rangle \\ &= \langle V_{\rho_{ii}}^* \Phi_{\rho_{ii}}(b^*a)V_{\rho_{ii}}\xi, \eta \rangle \\ &= \langle \Phi_{\rho_{ii}}(a)V_{\rho_{ii}}\xi, \Phi_{\rho_{ii}}(b)V_{\rho_{ii}}\eta \rangle \end{aligned}$$

for all $a, b \in A$ and for all $\xi, \eta \in E$, there is a unitary operator U_i from $E_{\rho_{ii}}$ to E_ρ such that

$$U_i(\Phi_{\rho_{ii}}(a)V_{\rho_{ii}}\xi) = \Phi_\rho(a)V_{\rho,i}\xi$$

[14, Theorem 3.5]. Moreover, $U_i\Phi_{\rho_{ii}}(a) = \Phi_\rho(a)U_i$ for all $a \in A$. Then $\widetilde{U_i}$, the unique extension of U_i to a bounded module homomorphism from $\widetilde{E_{\rho_{ii}}}$ to $\widetilde{E_\rho}$ with $\|U_i\| = \|\widetilde{U_i}\|$, is a unitary element in $\mathcal{L}_{B^{**}}(\widetilde{E_{\rho_{ii}}}, \widetilde{E_\rho})$.

Let $T \in [0, I]_\rho$. Then $\tilde{U}_i^* T \tilde{U}_i \in [0, I]_{\rho_{ii}}$, and since ρ_{ii} is pure, by Proposition 3.1, $\tilde{U}_i^* T \tilde{U}_i = \alpha I_{\widetilde{E_{\rho_{ii}}}}$ for some positive number α . Consequently, $T = \alpha I_{\widetilde{E_\rho}}$ and so ρ is pure. ■

In the following corollary we determine a class of extreme points in the set of all identity preserving completely positive linear maps from a unital locally C^* -algebra A to the C^* -algebra $M_n(B)$ of all $n \times n$ matrices with elements in the unital C^* -algebra B . This is a generalization of Corollaries 2.7 in [11] and 4.5 in [10].

COROLLARY 3.4 *Let A be a unital locally C^* -algebra, let B be a unital C^* -algebra and let $\rho = [\rho_{ij}]_{i,j=1}^n \in CP_\infty^n(A, B)$. If ρ_{ii} , $i \in \{1, \dots, n\}$ are unitarily equivalent pure unital completely positive linear maps from A to B and for all $i, j \in \{1, \dots, n\}$ with $i \neq j$, $\rho_{ij}(1) = 0$ and there is a unitary element u_{ij} in A such that $\rho_{ij}(u_{ij}) = 1$, then the map φ from A to $M_n(B)$ defined by $\varphi(a) = [\rho_{ij}(a)]_{i,j=1}^n$ is an extreme point in the set of all identity preserving completely positive linear maps from A to $M_n(B)$.*

Proof. Let φ_1 and φ_2 be two identity preserving completely positive linear maps from A to $M_n(B)$ and let $\alpha \in (0, 1)$ such that $\alpha\varphi_1 + (1 - \alpha)\varphi_2 = \varphi$. Then $\alpha\mathcal{S}^{-1}(\varphi_1) + (1 - \alpha)\mathcal{S}^{-1}(\varphi_2) = \rho$. From this relation and Propositions 3.3 and 3.1, we conclude that $\alpha\mathcal{S}^{-1}(\varphi_1) = \beta_1\rho$ for some positive number β_1 and $(1 - \alpha)\mathcal{S}^{-1}(\varphi_2) = \beta_2\rho$ for some positive number β_2 . Consequently, $\alpha\varphi_1 = \beta_1\varphi$ and $(1 - \alpha)\varphi_2 = \beta_2\varphi$. From these facts, since $\varphi_1(1) = \varphi_2(1) = \varphi(1) = I_n$, where I_n is the unity matrix in $M_n(B)$, we deduce that $\alpha = \beta_1$ and $1 - \alpha = \beta_2$. Therefore $\varphi_1 = \varphi_2 = \varphi$, and so φ is an extreme point in the set of all identity preserving completely positive linear maps from A to $M_n(B)$. ■

Let A be a unital locally C^* -algebra, let B be a C^* -algebra and let E be a Hilbert B -module. We denote by $CP_\infty^n(A, \mathcal{L}_B(E), I)$ the set of all completely n -positive linear maps $\rho = [\rho_{ij}]_{i,j=1}^n$ from A to $\mathcal{L}_B(E)$ such that $\rho_{ii}(1) = I_E$ for all $i \in \{1, \dots, n\}$ and $\rho_{ij}(1) = 0$ for all $i, j \in \{1, \dots, n\}$ with $i \neq j$.

The following theorem is a generalization of Theorems 3.8 in [18] and 4.6 in [10].

THEOREM 3.5 *Let $\rho \in CP_\infty^n(A, \mathcal{L}_B(E), I)$. Then ρ is an extreme point in the set $CP_\infty^n(A, \mathcal{L}_B(E), I)$ if and only if the map $T \rightarrow \left[\widetilde{V_{\rho,i}}^* T \widetilde{V_{\rho,j}} \right]_{i,j=1}^n$ from $C(\rho)$ to $M_n(\mathcal{L}_{B^{**}}(\widetilde{E}))$ is injective.*

Proof. Suppose that ρ is an extreme point in the set $CP_\infty^n(A, \mathcal{L}_B(E), I)$ and T is an element in $C(\rho)$ such that $\widetilde{V_{\rho,j}}^* T \widetilde{V_{\rho,i}} = 0$ for all $i, j \in \{1, \dots, n\}$. Since $\widetilde{V_{\rho,j}}^* T^* \widetilde{V_{\rho,i}} = \left(\widetilde{V_{\rho,i}}^* T \widetilde{V_{\rho,j}} \right)^*$ for all $i, j \in \{1, \dots, n\}$, we can suppose that $T = T^*$. It is not difficult to check that there are two positive numbers α and β such that $\frac{1}{4} I_{\widetilde{E_\rho}} \leq \alpha T + \beta I_{\widetilde{E_\rho}} \leq \frac{3}{4} I_{\widetilde{E_\rho}}$. Moreover, $\beta \in (0, 1)$. Let $T_1 = \frac{\alpha}{\beta} T + I_{\widetilde{E_\rho}}$ and $T_2 = I_{\widetilde{E_\rho}} - \frac{\alpha}{1-\beta} T$. Clearly, $T_k, k \in \{1, 2\}$ are two positive elements in $C(\rho)$. Then $\rho_{T_k} \in CP_\infty^n(A, \mathcal{L}_B(E))$, $k \in \{1, 2\}$ and since

$$(\rho_{T_k})_{ij}(1) = \widetilde{V_{\rho,i}}^* T_k \widetilde{V_{\rho,j}} \Big|_E = \widetilde{V_{\rho,i}}^* \widetilde{V_{\rho,j}} \Big|_E = V_{\rho,i}^* V_{\rho,j} = \begin{cases} I_E & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

for all $i, j \in \{1, \dots, n\}$ with $i \neq j$, and $k \in \{1, 2\}$, $\rho_{T_k} \in CP_\infty^n(A, \mathcal{L}_B(E), I)$ for each $k \in \{1, 2\}$. A simple calculus shows that $\beta \rho_{T_1} + (1 - \beta) \rho_{T_2} = \rho$, and since ρ is an extreme point, $\rho_{T_1} = \rho_{T_2} = \rho$. But $\rho_{T_1} = \frac{\alpha}{\beta} \rho_T + \rho$ and $\rho_{T_2} = \rho - \frac{\alpha}{1-\beta} \rho_T$. Therefore $\rho_T = 0$ and by Theorem 2.6, $T = 0$.

Conversely, suppose that the map $T \rightarrow \left[\widetilde{V_{\rho,i}}^* T \widetilde{V_{\rho,j}} \right]_{i,j=1}^n$ from $C(\rho)$ to $M_n(\mathcal{L}_{B^{**}}(\widetilde{E}))$ is injective. Let $\theta, \sigma \in CP_\infty^n(A, \mathcal{L}_B(E), I)$ and $\alpha \in (0, 1)$ such that $\alpha \theta + (1 - \alpha) \sigma = \rho$. By Theorem 2.6, there are two elements T_1 and T_2 in $[0, I]_\rho \subseteq C(\rho)$, such that $\alpha \theta = \rho_{T_1}$ and $(1 - \alpha) \sigma = \rho_{T_2}$. Then

$$\widetilde{V_{\rho,i}}^* T_1 \widetilde{V_{\rho,j}} \Big|_E = (\rho_{T_1})_{ij}(1) = \alpha \theta_{ij}(1) = \begin{cases} \alpha I_E & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

and

$$\widetilde{V_{\rho,i}}^* T_2 \widetilde{V_{\rho,j}} \Big|_E = (\rho_{T_2})_{ij}(1) = (1 - \alpha) \sigma_{ij}(1) = \begin{cases} (1 - \alpha) I_E & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}.$$

Therefore

$$\widetilde{V_{\rho,i}}^* \left(T_1 - \alpha I_{\widetilde{E_\rho}} \right) \widetilde{V_{\rho,j}} \Big|_E = \widetilde{V_{\rho,i}}^* \left(T_2 - (1 - \alpha) I_{\widetilde{E_\rho}} \right) \widetilde{V_{\rho,j}} \Big|_E = 0$$

from all $i, j \in \{1, \dots, n\}$. From these facts, Remark 1.4 and taking into account that $\widetilde{V_{\rho,i}}^* T_1 \widetilde{V_{\rho,j}} \Big|_E$ and $\widetilde{V_{\rho,i}}^* T_2 \widetilde{V_{\rho,j}} \Big|_E$ are elements in $\mathcal{L}_B(E)$ for all $i, j \in \{1, \dots, n\}$, we conclude that

$$\widetilde{V_{\rho,i}}^* \left(T_1 - \alpha I_{\widetilde{E_\rho}} \right) \widetilde{V_{\rho,j}} = \widetilde{V_{\rho,i}}^* \left(T_2 - (1 - \alpha) I_{\widetilde{E_\rho}} \right) \widetilde{V_{\rho,j}} = 0$$

for all $i, j \in \{1, \dots, n\}$. Hence $T_1 = \alpha I_{\widetilde{E_\rho}}$ and $T_2 = (1 - \alpha) I_{\widetilde{E_\rho}}$ and consequently $\theta = \sigma = \rho$. ■

References

- [1] W. ARVESON, *Subalgebras of C^* -algebras*, Acta Math., **123**(1969), 141-224.
- [2] S.J. BHATT AND D.J. KARIA, *Complete positivity and C^* -nuclearity for inverse limits of C^* -algebras*, Proc. Indian Acad. Sci. (Math. Sci.) **101** (1991), 149-167.
- [3] M. FRAGOULOPOULOU, *An introduction to the representation theory of topological $*$ algebras*, Schriftenreihe, Univ. Münster, **48** (1988), 1-81.
- [4] J. HEO, *Completely multi-positive linear maps and representations on Hilbert C^* -modules*, J. Operator Theory, **41**(1999), 3-22.
- [5] J. HEO, *A Radon-Nikodym theorem for completely positive invariant multilinear maps and its applications*, Math. Proc. Camb. Phil. Soc., **132**(2002), 181-192.
- [6] A. INOUE, *Locally C^* -algebras*, Mem. Faculty Sci. Kyushu Univ. Ser. A, **25**(1971), 197-235.
- [7] M. JOIȚA, *Strict completely positive maps between locally C^* -algebras and representations on Hilbert modules*, J. London Math. Soc. (2), **66**(2002), 421-432.
- [8] M. JOIȚA, *Multi-positive linear functionals on locally C^* -algebras*, Rend. Circ. Mat. Palermo (2), **53**(2004), 185-194.
- [9] M. JOIȚA, *Completely multi-positive linear maps between locally C^* -algebras and representations on Hilbert modules*, Studia Math.(to appear).
- [10] M. JOIȚA, *A Radon-Nikodym theorem for completely n -positive linear maps on locally C^* -algebras and its applications*, (submitted).
- [11] A. KAPLAN, *Multi-states on C^* -algebras*, Proc. Amer. Math. Soc., **106** (1989), 437-446.
- [12] I. KAPLANSKY, *Modules over operator algebras*, Amer. J. Math., **75**(1953), 839-853.
- [13] G.G.KASPAROV, *Hilbert C^* -modules: theorems of Stinespring and Voiculescu*, J. Operator Theory, **4**(1980),133-150.

- [14] E. C. LANCE, *Hilbert C^* -modules. A toolkit for operator algebraists*, London Mathematical Society Lecture Note Series 210, Cambridge University Press, Cambridge 1995.
- [15] H. LIN, *Bounded module maps and pure completely positive maps*, J. Operator Theory, **26**(1991), 121-139.
- [16] W. L. PASCHKE, *Inner product modules over B^* -algebras*, Trans. Amer. Math. Soc., **182**(1973), 443-468.
- [17] N. C. PHILLIPS, *Inverse limits of C^* -algebras*, J. Operator Theory, **19** (1988), 159-195.
- [18] W. STINESPRING, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc., **6**(1955), 211-216.
- [19] S. K. TSUI, *Completely positive module maps and completely positive extreme maps*, Proc. Amer. Math. Soc., **124**(1996), 437-445.

DEPARTMENT OF MATHEMATICS
 FACULTY OF CHEMISTRY
 UNIVERSITY OF BUCHAREST
 BD. REGINA ELISABETA NR.4-12
 BUCHAREST, ROMANIA
 mjoita@fmi.unibuc.ro